

VARIATIONAL LIMITS OF k -NN GRAPH-BASED FUNCTIONALS ON DATA CLOUDS

NICOLÁS GARCÍA TRILLOS

ABSTRACT. We consider i.i.d. samples $\mathbf{x}_1, \dots, \mathbf{x}_n$ from a measure ν with density supported on a bounded Euclidean domain $D \subseteq \mathbb{R}^d$ where $d \geq 3$. A graph on the point cloud is obtained by connecting two points if one of them is among the k -nearest neighbors of the other. Our goal is to study consistency of graph based procedures to clustering, classification and dimensionality reduction by studying the variational convergence of the graph total variation associated to such k -NN graph. We prove that provided $k := k_n$ scales like $n \gg k_n \gg \log(n)$, then the Γ -convergence of the graph total variation towards an appropriate weighted total variation is guaranteed.

1. INTRODUCTION

The development of machine learning and data analysis during the last decades has created new research opportunities for mathematicians and statisticians. Tasks like data clustering, data classification, and dimensionality reduction have been approached using methods whose statistical properties are of interest both for theoretical and practical reasons. In this paper we consider a broad class of approaches to such tasks known as *graph-based methods*, and in particular those that can be posed as variational problems defined on a graph. In general, the set of vertices on the graph is a data cloud $X_n := \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ and the graph structure comes from a similarity matrix for the points. Spectral clustering, total variation clustering, and diffusion maps, are all examples of graph-based methods with a variational flavor (see [5, 6, 16, 17, 21, 22, 26]).

In the setting where the data points are assumed to be draws from some unknown ground-truth distribution ν supported on $\mathbb{R}^d$, it is standard to use the Euclidean distance to induce the similarity among points in the cloud in such a way that points that are closer together are more similar than those that are further apart. The most popular graph constructions of this type are ε -graphs and k -NN (k nearest neighbors) graphs. For ε -graphs one needs to fix a decreasing kernel $\eta : [0, \infty) \rightarrow [0, \infty)$ and a length-scale $\varepsilon > 0$; in the literature the kernel is typically chosen to be a Gaussian kernel. The weight between two points $\mathbf{x}_i$ and $\mathbf{x}_j$ then takes the form

$$W_{ij} := \eta \left(\frac{|\mathbf{x}_i - \mathbf{x}_j|}{\varepsilon} \right).$$

k -NN graphs, on the other hand, are obtained by connecting two points with an edge if one of the points is among the k -nearest neighbors of the other. One can view the resulting graph as a weighted graph in which the weight between $\mathbf{x}_i$ and $\mathbf{x}_j$ is given by

$$(1.1) \quad J_k(\mathbf{x}_i, \mathbf{x}_j) := \begin{cases} 1 & \mathbf{x}_i \sim_k \mathbf{x}_j \\ 0 & \text{otherwise,} \end{cases}$$

where in the above and in the remainder of the paper we write $\mathbf{x}_i \sim_k \mathbf{x}_j$ whenever $\mathbf{x}_i$ is among the k -nearest neighbors of $\mathbf{x}_j$ or vice versa.

The goal of this paper is to establish rigorous stability results for variational problems defined on k -NN graphs that are closely connected to tasks like data clustering and data classification (see the examples in Section 2 below to get an idea of the possibilities); the variational problems of interest are defined in terms of a graph total variation energy or a graph Dirichlet energy (see Section 2 below for definitions). By establishing the stability of such variational problems we

deduce consistency results for a large variety of statistical procedures defined in terms of k -NN graphs.

It is important to highlight that in spite of the popularity of k -NN graphs among practitioners and certain advantages that they have over ε -graphs (see [26] for a discussion), the theoretical aspects of graph-based procedures have been mostly analyzed only in the context of ε -graphs. Indeed, asymptotic (as the sample size grows) properties of graph Laplacians have been studied by Belkin and Niyogi [4], Coifman and Lafon [6], Giné and Koltchinskii [13], Hein, Audibert and von Luxburg [15], and Singer [23]. These results deal mostly with pointwise consistency of graph Laplacians. The work of Arias-Castro, Pelletier and Pudlo [1] study the pointwise convergence of Cheeger energy and consequently of total variation, as well as variational convergence when the discrete functional is considered over an admissible set of characteristic functions which satisfy a “regularity” requirement. Spectral convergence of graph Laplacians, (relevant for data clustering) has been studied, among others, by Huang, and Jordan [25], Belkin and Niyogi [3], von Luxburg, Belkin and Bousquet [27], Singer and Wu [24]. Most of the results previously mentioned are obtained using tools from perturbation theory of linear operators.

A different set of tools were introduced in [9] and later used in [8, 11, 12]. The notion of Γ -convergence (see [7] and Section 2.1 below) and the introduction of a suitable metric (the TL^1 metric; see [9] and Section 2.1 below) allowing to compare objects at the discrete level with objects in the continuum, were crucial tools when establishing consistency results for a large class of balanced graph cuts in [12] and establishing consistency of spectral clustering under different choices of graph Laplacians in [11]. In all these results, sharp convergence rates for $\varepsilon := \varepsilon_n$ guaranteeing the consistency were provided; the assumptions on the kernel η were as general as possible.

Here we follow closely the framework from [9] and in particular our main results are analogues, in the context of k -NN graphs, of the results in [9] for ε -graphs.

1.1. Outline. The rest of the paper is organized as follows. In Section 2 we make our assumptions precise, present some of the examples of variational problems that are relevant for important tasks in data analysis, and lastly present the main results of the paper. In Section 2.1 we give the definitions of the TL^1 space and the notion of Γ -convergence; we also present some auxiliary results that are needed in the remainder. Finally, in Section 3 we present the proofs of the main results.

2. SET-UP AND MAIN RESULTS

Let $D \subseteq \mathbb{R}^d$ be a connected, bounded, open set with Lipschitz boundary. Let $\rho : D \rightarrow \mathbb{R}$ be a continuous density function bounded above and below by positive constants. Namely, assume that there are positive constants $0 < \rho_{\min} < \rho_{\max}$ such that for every $x \in D$ we have

$$(2.1) \quad \rho_{\min} \leq \rho(x) \leq \rho_{\max}, \quad \forall x \in D.$$

Let ν be the probability measure with density ρ , i.e., $d\nu(x) = \rho(x)dx$.

Let $X_n := \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be i.i.d samples from the distribution ν and denote by ν_n the empirical measure

$$\nu_n := \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{x}_i}.$$

Remark 2.1. Because ν has a density, with probability one, the condition $\mathbf{x}_i \sim_k \mathbf{x}_j$ is equivalent to

$$\nu_n(B(\mathbf{x}_i, r)) \leq \frac{k}{n} \quad \text{or} \quad \nu_n(B(\mathbf{x}_j, r)) \leq \frac{k}{n},$$

where $r = |\mathbf{x}_i - \mathbf{x}_j|$. Without the loss of generality we assume this fact in the remainder.

For $k := k_n \in \mathbb{N}$, we let $\bar{\varepsilon}_n$ be the number for which

$$(2.2) \quad \bar{\varepsilon}_n^d := \frac{k_n}{n}.$$

We define the *graph total variation* functional as

$$GTV_{n,k}(u_n) := \frac{1}{n^2 \varepsilon_n^{d+1}} \sum_{i,j} J_k(\mathbf{x}_i, \mathbf{x}_j) |u_n(\mathbf{x}_i) - u_n(\mathbf{x}_j)|, \quad u_n \in L^1(\nu_n).$$

We are interested in minimization problems whose objective functions involve the energy $GTV_{n,k}$. Let us list some examples that are relevant for clustering, classification and dimensionality reduction.

Example 2.2. (Cheeger and ratio graph cuts for clustering). Consider a weighted graph with vertices $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ and weight matrix $(w_{ij})_{i,j}$. The *Cheeger cut* functional associated to this graph is defined by

$$Cut_C(A, A^c) := \frac{\sum_{\mathbf{x}_i \in A} \sum_{\mathbf{x}_j \notin A} w_{ij}}{\min\{|A|, |A^c|\}}, \quad A \subseteq X_n,$$

where $|A|$ and $|A^c|$ are the number of vertices in A and A^c respectively. A similar functional is the *ratio cut* defined by

$$Cut_R(A, A^c) := \frac{\sum_{\mathbf{x}_i \in A} \sum_{\mathbf{x}_j \notin A} w_{ij}}{|A| \cdot |A^c|}, \quad A \subseteq X_n.$$

Both of the above functionals can be seen as examples of a larger class of functionals known as *balanced cuts*. These type of functionals penalize partitions $\{A, A^c\}$ of X_n that either have a big interface separating A and A^c or are highly unbalanced in the sense that A and A^c are of dissimilar size; intuitively, these are desirable features for a good partitioning of the data and motivate considering the minimization problem

$$\min_{A \subseteq X_n} Cut(A, A^c),$$

to obtain a good partition of X_n . Notice that the denominator in both functionals is, up to a multiplicative factor, the graph total variation defined by

$$GTV(\mathbb{1}_A) := \sum_{i,j} w_{ij} |\mathbb{1}_A(\mathbf{x}_i) - \mathbb{1}_A(\mathbf{x}_j)|.$$

Assuming that the set of vertices consists of samples from the distribution ν and that the weights of the graph are those obtained from an ε -graph (with $\varepsilon := \varepsilon_n$ chosen appropriately), the results in [12] state that minimizers of the Cheeger and ratio cuts converge, as $n \rightarrow \infty$, to objects in the continuum with analogue properties to those of the minimizers at the discrete level. The limit (taken in the TL^1 sense; a suitable strong topology) is found to be a minimizer of an analogue continuous Cheeger/ratio cut (see [12] for more details). This result can then be interpreted as a consistency result for clustering using balanced cuts.

Example 2.3. (Graph total variation in the context of classification). Suppose that

$$\{(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_n, y_n)\}$$

are samples from some distribution ν supported on $\overline{D} \times \{0, 1\}$. The pair $(\mathbf{x}_i, y_i)$ is interpreted as the feature values ($\mathbf{x}$ variable) and label (y variable) of an individual from a given population. Based on the observed data (training data) the idea is to construct a “good” *classifier* assigning labels to every potential individual in the population.

A typical choice of risk functional used to define “good” classifiers is the *average missclassification error*. Unfortunately, when the distribution ν is unknown (as it is usually the case) it is not possible to determine the classifier that minimizes the average misclassification error (Bayes classifier) and hence an approximation of it based exclusively on the observed data is the best thing one can hope for. Such an approximation can be obtained using the graph total variation as we describe below.

Given the weighted graph $(\{\mathbf{x}_i\}_i, (w_{ij})_{i,j})$, consider the energy

$$R_{n,\lambda}(u) := R_n(u) + \lambda GTV(u), \quad u : \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \rightarrow \mathbb{R}.$$

The functional R_n is the *empirical risk* and the parameter λ is introduced so as to emphasize or deemphasize the regularizing effect of the graph total variation. In [8], the weights for the graph are assumed to be those coming from an ε -graph and the problem of obtaining an approximation to the Bayes classifier is divided into first solving the minimization problem

$$\min_{u: X_n \rightarrow \mathbb{R}} R_{n,\lambda}(u),$$

and then extend the minimizer to the whole ambient space appropriately. With the right scaling for $\lambda = \lambda_n$, one can establish the asymptotic consistency of the constructed approximation. See [8] for more details.

Example 2.4. (Spectral clustering and spectral embeddings). Undoubtedly, one of the most popular graph based methods for data clustering is *spectral clustering*. In the two way clustering setting, spectral clustering can be seen as a relaxation of the ratio cut minimization problem mentioned in Example 2.2. Indeed, the relaxed problem takes the form

$$\min_{u: X_n \rightarrow \mathbb{R}} \frac{\sum_{i,j} w_{ij} (u(\mathbf{x}_i) - u(\mathbf{x}_j))^2}{\sum_i (u(\mathbf{x}_i) - \bar{u})^2},$$

where $\bar{u}$ is the average $\bar{u} := \frac{1}{n} \sum_i u(v_i)$. It is well known that the above optimization problem is actually an eigenvalue problem for the graph Laplacian and that the first non-trivial eigenvector of the graph Laplacian is a minimizer.

In the context of multi-way clustering, higher eigenvectors of the graph Laplacian are used to define an embedding of the data cloud into a Euclidean space with low dimension. In turn, the embedded data can be clustered using an algorithm like k -means, inducing in this way a partition for the original data.

Among the many results in the literature addressing the consistency of spectral clustering (see the Introduction for some references), we highlight the work in [11] which exploits the variational characterization of eigenvectors/eigenvalues. The graph on the cloud X_n is assumed to be an ε -graph.

Our goal is to study the limiting behavior, as the sample size goes to infinity, of minimizers of optimization problems like the ones presented in the above examples when the graph considered is a k -NN graph and $k = k_n$ scales with the sample size in a suitable way. As explained in [8, 9, 11, 12] in the context of ε -graphs, most of the convergence analysis depends on establishing the variational convergence (Γ -convergence) of the graph total variation towards an analogue continuous total variation; the TL^1 distance introduced in [9] was suitable for this purpose. We focus on establishing the Γ -limit (in the TL^1 -sense) of the functionals GTV_{n,k_n} for an appropriate scaling $k = k_n$. The limiting functional turns out to be a weighted total variation of the type that we describe below.

Definition 2.5. Let $h : D \rightarrow \mathbb{R}$ be a continuous function bounded below and above by positive constants. For an arbitrary function $u \in L^1(D)$, we define its weighted (by h) total variation as

$$(2.3) \quad TV(u; h) := \sup \left\{ \int_D u(x) \operatorname{div}(\zeta) dx : \zeta \in C_c^\infty(D; \mathbb{R}^d), \quad |\zeta(x)| \leq h(x) \quad \forall x \in D \right\}.$$

In the above and in the remainder of the paper, we use $L^1(D)$ to represent the space of L^1 functions with respect to the Lebesgue measure restricted to D . In addition, when $h \equiv 1$ we write $TV(u)$ instead of $TV(u; h)$ and we denote by $BV(D)$ the space of functions $u \in L^1(D)$ for which $TV(u) < \infty$. Notice that for any h continuous and bounded above and below by positive constants, the condition $TV(u) < \infty$ is equivalent to the condition $TV(u; h) < \infty$.

Remark 2.6. If $u \in BV(D)$ is smooth, then

$$TV(u; h) = \int_D |\nabla u| h(x) dx.$$

Also, if $u = \mathbb{1}_A$ for some open set A with smooth boundary, then

$$TV(u; h) = \int_{\partial A \cap D} h(x) d\mathcal{H}^{d-1}(x),$$

where $\mathcal{H}^{d-1}$ is the $(d-1)$ -dimensional Hausdorff measure.

Before we state our main results, let us introduce some functions that we use in the remainder. Let $\eta : [0, \infty) \rightarrow \mathbb{R}$ be the Heaviside function given by

$$\eta(t) := \begin{cases} 1 & \text{if } t < 1 \\ 0 & \text{if } t \geq 1, \end{cases}$$

For an arbitrary $\varepsilon > 0$ we let η_ε be the function

$$\eta_\varepsilon(t) := \frac{1}{\varepsilon^d} \eta\left(\frac{t}{\varepsilon}\right), \quad t \in [0, \infty).$$

Finally, we let σ_η be the quantity

$$(2.4) \quad \sigma_\eta := \int_{\mathbb{R}^d} \eta(z) |z_1| dz,$$

where z_1 is the first coordinate of the vector z . Notice that in the above expression z_1 can be replaced with $z \cdot v$ for any vector v with unit norm.

The main results of this paper are as follows.

Theorem 2.7. (*Γ -convergence*) Let $d \geq 3$ and let $D \subseteq \mathbb{R}^d$ be an open, connected, bounded domain with Lipschitz boundary. Let $\rho : D \rightarrow \mathbb{R}$ be a continuous density function satisfying (2.1). Let $\{k_n\}_{n \in \mathbb{N}}$ be a sequence of natural numbers satisfying

$$\lim_{n \rightarrow \infty} \frac{k_n}{\log(n)} = +\infty, \quad \lim_{n \rightarrow \infty} \frac{k_n}{n} = 0.$$

Then, the functionals GTV_{n, k_n} Γ -converge towards $\frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(\cdot; \rho^{1-1/d})$ in the TL^1 -sense, where σ_η is defined in (2.4), and α_d is the volume of the unit ball in $\mathbb{R}^d$. Furthermore, when restricted to indicator functions, the energies GTV_{n, k_n} Γ -converge to the functional $\frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(\cdot; \rho^{1-1/d})$ restricted to indicator functions.

Theorem 2.8. (*Compactness*) Under the same assumptions in Theorem 2.7, the following statement holds with probability one: If $\{u_n\}_{n \in \mathbb{N}}$ is a sequence with $u_n \in L^1(\nu_n)$ for which

$$\sup_{n \rightarrow \infty} \|u_n\|_{L^1(\nu_n)} < \infty$$

and

$$\sup_{n \rightarrow \infty} GTV_{n, k_n}(u_n) < \infty,$$

then $\{u_n\}_{n \in \mathbb{N}}$ is pre-compact in $TL^1(D)$. That is, every subsequence of $\{u_n\}_{n \in \mathbb{N}}$ has a further subsequence converging in $TL^1(D)$.

Remark 2.9. The above theorems hold for domains D in $\mathbb{R}^2$ provided we replace the condition $\lim_{n \rightarrow \infty} \frac{k_n}{\log(n)} = \infty$ with the condition $\lim_{n \rightarrow \infty} \frac{k_n}{(\log(n))^{3/2}} = \infty$. This can be seen directly from our proofs, and follows from the fact that the rate of convergence of the ∞ -transportation distance between ν and ν_n (for $d = 2$) scales like $\frac{\log(n)^{3/4}}{n^{1/2}}$ and not like $\frac{(\log(n))^{1/2}}{n^{1/2}}$ (see [10]).

Remark 2.10. The above set-up and results can be extended to the setting in which the support of ν is not a domain D contained in the ambient space $\mathbb{R}^d$ but a compact manifold $\mathcal{M} \subseteq \mathbb{R}^d$ with intrinsic dimension m . When that is the case, the appropriate scalings for the functionals are obtained using the intrinsic dimension m and not the dimension of the ambient space d . Although in this paper we omit the details of such extension, we point out that the results in [20] (later used in the proofs in [9]) can be adapted to the manifold setting by establishing results

analogues to those in [10] in the manifold setting, using the geodesic flow in $\mathcal{M}$, and the fact that the Euclidean distance (distance in the ambient space $\mathbb{R}^d$) is a third order approximation for the geodesic distance (intrinsic distance), i.e. ,

$$d_{\mathcal{M}}(x, y) = |x - y| + O(|x - y|^3), \quad x, y \in \mathcal{M}.$$

The details can be presented elsewhere in a more general setting which is also of interest; see Remark 2.11 below.

Remark 2.11. Suppose that $d \geq 3$. One of the important consequences of Theorem 2.7 is that the admissible regimes for $k = k_n$ that guarantee the recovery of a non-trivial variational limit for the energies $GTV_{n,k}$ does not depend on d (we impose $n \gg k_n \gg \log(n)$). As a consequence, notice that despite the fact that the scaling factor appearing in $GTV_{n,k}$ depends on d , the minimizers of the Cheeger and ratio cuts are independent of such scaling given that the *argmin* is invariant under the rescaling of the energy by a positive constant. In other words, in principle we do not need to know the dimensionality of the data before hand in order to obtain good clusters using the graph total variation associated to a k -NN graph. Notice that the same is not true for ε -graphs.

In particular, it is plausible to extend the consistency of Cheeger and ratio cuts results to the setting in which the dimensionality of the data changes in space, i.e., if ν 's support is something like a union of manifolds with different intrinsic dimensions. This problem will be explored in the future.

Remark 2.12. Although we have restricted our attention to the graph total variation $GTV_{n,k}$, it is possible to obtain analogues to Theorem 2.7 and Theorem 2.8 for the graph Dirichlet energy

$$GD_{n,k}(u_n) := \frac{1}{n^2 \varepsilon_n^{d+2}} \sum_{i,j} J_k(\mathbf{x}_i, \mathbf{x}_j) (u_n(\mathbf{x}_i) - u_n(\mathbf{x}_j))^2, \quad u_n \in L^2(\nu_n).$$

The results in [11] for ε -graphs can then be adapted to the k -NN graph case. As mentioned in Example 2.4, the functional $GD_{n,k}$ is of relevance for spectral clustering.

2.1. Preliminaries. The purpose of this subsection is to present some definitions and preliminary results that we use in the proof of our main theorems. In particular we present the definitions of TL^1 -space and Γ -convergence.

$TL^1(D)$ is defined as the set of all pairs (μ, u) where μ is a Borel probability measure on D and u is a function in $L^1(D, \mu)$ (written $L^1(\mu)$ from now on). This set can be endowed with the TL^1 -metric defined by

$$d_{TL^1}((\mu_1, u_1), (\mu_2, u_2)) := \inf_{\pi \in \Gamma(\mu_1, \mu_2)} \left\{ \int_{D \times D} |x - y| d\pi(x, y) + \int_{D \times D} |u_1(x) - u_2(y)| d\pi(x, y) \right\},$$

where $\Gamma(\mu_1, \mu_2)$ stands for the set of couplings between μ_1 and μ_2 , that is, the set of all probability measures on the product $D \times D$ whose first and second marginals are μ_1 and μ_2 respectively.

Given that with probability one the empirical measure ν_n converges weakly to ν as $n \rightarrow \infty$, and given that ν has a density, we may use the characterization of TL^1 -convergence from Proposition 3.12 in [9] to conclude that $\{(\nu_n, u_n)\}_{n \in \mathbb{N}}$ converges to (ν, u) in TL^1 if and only if *there exists* a sequence of transportation maps $\{T_n\}_{n \in \mathbb{N}}$ with $T_{n\#}\nu = \nu_n$ (T_n pushes forward ν into ν_n), satisfying

$$(2.5) \quad \lim_{n \rightarrow \infty} \int_D |T_n(x) - x| d\nu(x) = 0$$

and

$$(2.6) \quad \lim_{n \rightarrow \infty} \int_D |u_n(T_n(x)) - u(x)| d\nu(x) = 0.$$

In turn, this holds if and only if *for all* sequences of transportation maps with $T_{n\#}\nu = \nu_n$ that satisfy (2.5) one has (2.6). Because of this characterization, we may abuse notation slightly and

simply write $u_n \xrightarrow{TL^1} u$ without specifying the corresponding attached measures whenever it is clear from context that they can be omitted.

A special choice of transportation maps between ν and ν_n that we use in the remainder are provided by [10]. We may consider transportation maps $T_n : D \rightarrow X_n$ between the measures ν and ν_n satisfying the condition

$$(2.7) \quad \|Id - T_n\|_\infty \leq \frac{C(\log(n))^{1/d}}{n^{1/d}} =: \delta_n,$$

for some constant C (provided $d \geq 3$). Indeed, the results in [10] show that with probability one, there exists transportation maps $\{T_n\}_{n \in \mathbb{N}}$ for which $T_{n\#}\nu = \nu_n$ and for which (2.7) holds for all large enough $n \in \mathbb{N}$. Since all of our results are asymptotic in nature, we may as well assume for the remainder that with probability one (2.7) holds for all $n \in \mathbb{N}$. One last relevant property of the map T_n , which follows directly from the fact that it transports ν into ν_n , is the change of variables formula

$$(2.8) \quad \int_D f(x) d\nu_n(x) = \int_D f(T_n(x)) d\nu(x),$$

which allows us to write integrals with respect to ν_n in terms of integrals with respect to ν .

We now present the definition of Γ -convergence in the context of a general metric space. This is a notion of convergence for functionals which together with a coercivity assumption, guarantee the stability of minimizers in the limit. A standard reference for Γ -convergence is [7].

Definition 2.13. *Let $(\mathcal{X}, d_{\mathcal{X}})$ be a metric space and let $F_n : \mathcal{X} \rightarrow [0, \infty]$ be a sequence of functionals. The sequence $\{F_n\}_{n \in \mathbb{N}}$ Γ -converges with respect to the metric $d_{\mathcal{X}}$ to the functional $F : \mathcal{X} \rightarrow [0, \infty]$ as $n \rightarrow \infty$ if the following inequalities hold:*

- **Liminf inequality:** *For every $x \in \mathcal{X}$ and every sequence $\{x_n\}_{n \in \mathbb{N}}$ converging to x ,*

$$\liminf_{n \rightarrow \infty} F_n(x_n) \geq F(x),$$

- **Limsup inequality:** *For every $x \in \mathcal{X}$ there exists a sequence $\{x_n\}_{n \in \mathbb{N}}$ converging to x satisfying*

$$\limsup_{n \rightarrow \infty} F_n(x_n) \leq F(x).$$

We say that F is the Γ -limit of the sequence of functionals $\{F_n\}_{n \in \mathbb{N}}$ (with respect to the metric $d_{\mathcal{X}}$).

Remark 2.14. A sequence $\{x_n\}_{n \in \mathbb{N}} \subseteq \mathcal{X}$ like the one appearing in the limsup inequality is said to be a recovery sequence for x . It is straightforward to show that if one can find recovery sequences for all elements in a set $\mathcal{X}'$ satisfying:

- (1) For all $x \in \mathcal{X}$ there exists a sequence $\{x_l\}_{l \in \mathbb{N}} \subseteq \mathcal{X}'$ such that $x_l \rightarrow x$ and $F(x_l) \rightarrow F(x)$, then one can find recovery sequences for all elements in $\mathcal{X}$; a set $\mathcal{X}'$ satisfying (1) above is said to be dense in $\mathcal{X}$ with respect to F . This fact follows from a simple diagonal argument (see [7]).

We have presented the notion of Γ convergence in this generality because some of the Γ -limits we will consider are taken in the context of the metric space $L^1(D)$ whereas others are taken in the context of the metric space $TL^1(D)$. Also, notice that when the functionals are allowed to be random (as it is the case for the graph total variation), Γ -convergence has to be interpreted as in the Definition 2.11 in [9].

To conclude this section we list two important properties of the weighted total variation functional $TV(\cdot; h)$. Let $h : D \rightarrow \mathbb{R}$ be a continuous function which is bounded above and below by positive constants. The first property is a representation formula for $TV(u; h)$ in terms of the distributional derivative of u . Indeed, it follows from the work in [2] that for every $u \in BV(D)$ one can write

$$(2.9) \quad TV(u; h) = \int_D h(x) d|Du|(x),$$

where in the above Du stands for the distributional derivative of u (which in general is a signed measure) and $|Du|$ stands for the total variation measure associated to Du .

The second property that is relevant for our purposes is the *coarea formula* which states that for every $u \in BV(D)$,

$$(2.10) \quad TV(u; h) = \int_{-\infty}^{\infty} TV(\mathbb{1}_{\{x : u(x) > t\}}; h) dt.$$

This formula says that the total variation of u can be written in terms of the total variation of its level sets. See Theorem 13.25 in [18].

3. PROOFS OF THEOREMS

The main auxiliary result that we need to establish our main results is the following.

Proposition 3.1. *For every $n \in \mathbb{N}$ let $\varepsilon_n, \hat{\varepsilon}_n : D \rightarrow (0, \infty)$ be functions satisfying*

$$\sup_{x \in D} \left| \frac{\varepsilon_n(x)}{\hat{\varepsilon}_n(x)} - 1 \right| \rightarrow 0 \quad , \quad \text{as } n \rightarrow \infty,$$

and

$$\sup_{x \in D} \varepsilon_n(x) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

For $n \in \mathbb{N}$ consider the energy $F_n : L^1(D) \rightarrow [0, \infty)$ defined by

$$F_n(u) := \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(x-y) |u(x) - u(y)| \rho(y) dy \right) \rho(x) dx, \quad u \in L^1(\nu),$$

where

$$\hat{f}_n(x) := \frac{(\hat{\varepsilon}_n(x))^d}{(\nu(B(x, \varepsilon_n(x))))^{1+1/d}}.$$

Then, as $n \rightarrow \infty$, F_n Γ -converges in the $L^1(D)$ sense towards F , where F is given by

$$F(u) := \frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(u; \rho^{1-1/d}).$$

Proposition 3.1 is a consequence of the next lemma.

Lemma 3.2. *Let $\rho_1, \rho_2 : \mathbb{R}^d \rightarrow (0, \infty)$ be two Lipschitz continuous functions which are bounded above and below by positive constants in D . Let μ be the Borel measure on $\mathbb{R}^d$ given by $d\mu(x) = \rho_2(x)dx$. Let $\varepsilon_n, \hat{\varepsilon}_n : D \rightarrow (0, \infty)$ be functions satisfying*

$$\sup_{x \in D} \left| \frac{\varepsilon_n(x)}{\hat{\varepsilon}_n(x)} - 1 \right| \rightarrow 0 \quad , \quad \text{as } n \rightarrow \infty,$$

and

$$\sup_{x \in D} \varepsilon_n(x) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Let $F_n : L^1(D) \rightarrow [0, \infty)$ be defined by

$$F_n(u) := \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(x-y) |u(x) - u(y)| \rho_1(y) dy \right) \rho_1(x) dx, \quad u \in L^1(\nu),$$

where

$$\hat{f}_n(x) := \frac{(\hat{\varepsilon}_n(x))^d}{(\mu(B(x, \varepsilon_n(x))))^{1+1/d}}.$$

Then, as $n \rightarrow \infty$, F_n Γ -converges in the $L^1(D)$ sense towards

$$F(u) := \frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV \left(u; \frac{\rho_1^2}{\rho_2^{1+1/d}} \right).$$

Proof. Let us start by proving the liminf inequality. That is, let us prove that for every $u \in L^1(D)$ and every sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq L^1(D)$ satisfying $u_n \xrightarrow{L^1(D)} u$, we have

$$(3.1) \quad \liminf_{n \rightarrow \infty} F_n(u_n) \geq F(u).$$

The following simplifications are standard. First, working along subsequences we may assume without the loss of generality that the liminf is actually a limit. In addition, we may assume that both of the terms involved in inequality 3.1 are finite. We now split the proof of (3.1) into several steps.

Step 1: Instead of working with the energy F_n directly, we first consider a simpler related energy E_n defined by

$$E_n(v) := \int_D \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_D \eta_{\hat{\varepsilon}_n(x)}(|x-y|)|v(x)-v(y)|dy \right) g(x)dx, \quad v \in L^1(D),$$

where

$$g(x) := \frac{(\rho_1(x))^2}{(\alpha_d \rho_2(x))^{1+1/d}}, \quad x \in D.$$

Notice that for every $x \in D$ we have

$$\left| \mu(B(x, \hat{\varepsilon}_n(x))) - \alpha_d \rho_2(x)(\hat{\varepsilon}_n(x))^d \right| \leq \alpha_d \text{Lip}(\rho_2)(\hat{\varepsilon}_n(x))^{d+1}.$$

From the previous inequality, the Lipschitz continuity of ρ_1, ρ_2 and the assumptions on $\hat{\varepsilon}_n(\cdot)$ and $\varepsilon_n(\cdot)$, we deduce that for every $x \in D$ and $y \in B(x, \hat{\varepsilon}_n(x))$,

$$\left| \frac{g(x)}{\hat{\varepsilon}_n(x)} - \hat{f}_n(x) \rho_1(x) \rho_1(y) \right| \leq C \sup_{z \in D} \varepsilon_n(z),$$

where C is a constant that does not depend on x or y . It then follows that

$$\lim_{n \rightarrow \infty} |F_n(u_n) - E_n(u_n)| = 0.$$

In particular, to obtain (3.1) we may as well show that

$$\liminf_{n \rightarrow \infty} E_n(u_n) \geq F(u).$$

Step 2: Let B be a closed ball contained in D . We claim that

$$\liminf_{n \rightarrow \infty} \int_B \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_B \eta_{\hat{\varepsilon}_n(x)}(|x-y|)|u_n(x)-u_n(y)|dy \right) dx \geq \sigma_\eta TV_B(u),$$

where $TV_B(u)$ is defined as

$$(3.2) \quad TV_B(u) := \sup \left\{ \int_{B^\circ} u(x) \text{div}(\zeta) dx : \zeta \in C_c^\infty(B^\circ; \mathbb{R}^d), \quad |\zeta(x)| \leq 1 \quad \forall x \in B^\circ \right\},$$

where in the above B° stands for the interior of the closed ball B (the ball without its boundary).

The claim follows from Theorem 8 in [20]. The only difference in the setting we consider here and the setting considered in [20] is that in our case the length-scale $\hat{\varepsilon}_n$ depends on location $\hat{\varepsilon}_n := \hat{\varepsilon}_n(x)$. The fact that $\hat{\varepsilon}_n(x)$ converges to zero uniformly over $x \in D$ as $n \rightarrow \infty$ is enough to make the proof in [20] carry through with essentially no modifications. We remark that the arguments in [20] were also used in [9]; in [9] the presence of a non-constant density makes computations a bit more tedious.

Step 3: Suppose that $u \in BV(D)$ is of the form $u = \mathbb{1}_A$ for some measurable set $A \subseteq D$. That is, suppose that u is the indicator function of a set with finite perimeter (with respect to D). We claim that there exists a sequence $\{\mathcal{F}_l\}_{l \in \mathbb{N}}$ of collections of closed balls contained in D with the following properties:

- (1) For every $l \in \mathbb{N}$, the family $\mathcal{F}_l$ is finite.
- (2) For every $l \in \mathbb{N}$, the balls $B := \overline{B(x_B, r_B)}$ in $\mathcal{F}_l$ are pairwise disjoint.
- (3) $\lim_{l \rightarrow \infty} \max\{r_B : B \in \mathcal{F}_l\} = 0$.
- (4) For every $l \in \mathbb{N}$, and for every $B \in \mathcal{F}_l$, $|Du|(\partial B) = 0$.

$$(5) \lim_{l \rightarrow \infty} |Du| \left(\bigcup_{B \in \mathcal{F}_l} B \right) = |Du|(D).$$

To prove this claim, let us start by noting that because A has finite perimeter, we may talk about its *reduced boundary* $\partial^* A$ (see [19]). In particular, for every measurable set $S \subseteq D$,

$$|Du|(S) = \mathcal{H}^{d-1}(\partial^* A \cap S),$$

where $\mathcal{H}^{d-1}$ stands for the $(d-1)$ -dimensional Hausdorff measure.

De Giorgi's structure theorem (see Theorem 15.9 in [19]) implies that there exist countably many C^1 -hypersurfaces $\{M_i\}_{i \in \mathbb{N}}$, compact sets $K_i \subseteq M_i$, and a $\mathcal{H}^{d-1}$ -null set F such that

$$\partial^* A = F \cup \bigcup_{i \in \mathbb{N}} K_i.$$

Fix $j \in \mathbb{N}$. From the fact that the sets K_i are subsets of C^1 -hypersurfaces, it is straightforward to see that one can find a sequence $\{\tilde{\mathcal{F}}_l^j\}_{l \in \mathbb{N}}$ of families of closed balls contained in D , satisfying (1), (2), (3) as above and such that

$$\lim_{l \rightarrow \infty} \mathcal{H}^{d-1} \left(\bigcup_{B \in \tilde{\mathcal{F}}_l^j} B \cap \bigcup_{i=1}^j K_i \right) = \mathcal{H}^{d-1} \left(D \cap \bigcup_{i=1}^j K_i \right).$$

Still working with j fixed, we now proceed to modify the balls in the families $\tilde{\mathcal{F}}_l^j$ to obtain balls that also satisfy (4). Indeed, we consider the family $\mathcal{F}_l^j$ of closed balls obtained from the balls in $\tilde{\mathcal{F}}_l^j$ by adding δ_l units to their original radius. The number δ_l can be chosen in such a way that $0 < \delta_l < \frac{1}{l}$, the balls in $\mathcal{F}_l^j$ are all contained in D , are pairwise disjoint and satisfy property (4). This can be done because $\partial^* A \cap D$ has finite $\mathcal{H}^{d-1}$ -measure and hence its intersection with the boundary of arbitrary balls centered at a fixed point, can only have non-zero $\mathcal{H}^{d-1}$ -measure for at most countably many of such balls.

A sequence of families $\{\mathcal{F}_l\}_{l \in \mathbb{N}}$ of balls with the desired properties can then be obtained from the families $\{\mathcal{F}_l^j\}_{l \in \mathbb{N}}$ for every $j \in \mathbb{N}$, by using a diagonal argument knowing that

$$\lim_{j \rightarrow \infty} \mathcal{H}^{d-1} \left(D \cap \bigcup_{i=1}^j K_i \right) = \mathcal{H}^{d-1}(D \cap \partial^* A).$$

With the families $\{\mathcal{F}_l\}_{l \in \mathbb{N}}$ at hand, we may now consider the functions

$$g_l(x) := \begin{cases} \min_{y \in B} g(y), & \text{if } x \in B, \quad B \in \mathcal{F}_l \\ 0, & \text{otherwise.} \end{cases}$$

Then, for every $l \in \mathbb{N}$

$$\begin{aligned} \liminf_{n \rightarrow \infty} E_n(u_n) &\geq \liminf_{n \rightarrow \infty} \sum_{B \in \mathcal{F}_l} \int_B \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_B \eta_{\hat{\varepsilon}_n(x)}(|x-y|) |u_n(x) - u_n(y)| dy \right) g(x) dx \\ &\geq \sum_{B \in \mathcal{F}_l} \liminf_{n \rightarrow \infty} \int_B \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_B \eta_{\hat{\varepsilon}_n(x)}(|x-y|) |u_n(x) - u_n(y)| dy \right) g(x) dx \\ &\geq \sum_{B \in \mathcal{F}_l} g_l(x_B) \liminf_{n \rightarrow \infty} \int_B \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_B \eta_{\hat{\varepsilon}_n(x)}(|x-y|) |u_n(x) - u_n(y)| dy \right) dx \\ &\geq \sum_{B \in \mathcal{F}_l} g_l(x_B) \sigma_\eta TV_B(u) \\ &= \sum_{B \in \mathcal{F}_l} g_l(x_B) \sigma_\eta \int_B d|Du|(x) \\ &= \sigma_\eta \int_D g_l(x) d|Du|(x), \end{aligned}$$

where in the fourth inequality we used Step 2 and in the first equality we used Lemma 15.12 in [19] combined with property (4) for the balls in $\mathcal{F}_l$.

Finally, from properties (3) and (5) of $\{\mathcal{F}_l\}_{l \in \mathbb{N}}$ and from the Lipschitz continuity of g we know that

$$\lim_{l \rightarrow \infty} g_l(x) = g(x), \quad \text{for } |Du| \text{ a.e. } x \in D.$$

The dominated convergence theorem implies that

$$\lim_{l \rightarrow \infty} \int_D g_l(x) d|Du|(x) = \int_D g(x) d|Du|(x) = TV(u; g),$$

where in the last equality we have used the representation formula (2.9). We conclude that for functions $u = \mathbb{1}_A \in BV(D)$ (i.e. indicator functions for sets of finite perimeter) inequality (3.1) holds.

Step 4: In order to prove (3.1) for general $u \in BV(D)$ we use Step 3 and the coarea formula for the energies E_n and F . Notice that the coarea formula for F is given in (2.10), whereas the coarea formula for E_n follows directly from the identity

$$|a - b| = \int_{-\infty}^{\infty} |\mathbb{1}_{a>t} - \mathbb{1}_{b>t}| dt, \quad \forall a, b \in \mathbb{R}.$$

If $u_n \xrightarrow{L^1(D)} u$, then for Lebesgue-a.e. $t \in \mathbb{R}$ we have

$$\mathbb{1}_{\{u_n > t\}} \xrightarrow{L^1(D)} \mathbb{1}_{\{u > t\}}, \quad \text{as } n \rightarrow \infty.$$

We may then apply Step 3 to conclude that for almost every $t \in \mathbb{R}$

$$\liminf_{n \rightarrow \infty} E_n(\mathbb{1}_{\{u_n > t\}}) \geq F(\mathbb{1}_{\{u > t\}}).$$

Integrating the above inequality with respect to t , and using Fatou's lemma and the coarea formulas for the energies E_n and F , we deduce that

$$\liminf_{n \rightarrow \infty} E_n(u_n) = \liminf_{n \rightarrow \infty} \int_{-\infty}^{\infty} E_n(\mathbb{1}_{\{u_n > t\}}) dt \geq \int_{-\infty}^{\infty} \liminf_{n \rightarrow \infty} E_n(\mathbb{1}_{\{u_n > t\}}) dt \geq \int_{-\infty}^{\infty} E(\mathbb{1}_{\{u > t\}}) dt = F(u).$$

Having established the liminf inequality we now proceed to establish the limsup inequality. From remark 2.14 and the density of $C_c^\infty(\mathbb{R}^d)$ functions in $L^1(D)$ with respect to TV (see Proposition 2.4 in [9]), it is enough to find a recovery sequence for every u obtained as the restriction to D of a function in $C_c^\infty(\mathbb{R}^d)$. We actually show that for every $u \in C_c^\infty(\mathbb{R}^d)$

$$(3.3) \quad \limsup_{n \rightarrow \infty} F_n(u) \leq F(u).$$

In other words $u_n := u$ for all n , is a recovery sequence for u . To prove this, notice that

$$\begin{aligned} E_n(u) &= \int_D \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_{B(x, \hat{\varepsilon}_n(x)) \cap D} \eta_{\hat{\varepsilon}_n(x)}(x - y) |u(x) - u(y)| dy \right) g(x) dx \\ &\leq \int_D \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_{B(x, \hat{\varepsilon}_n(x))} \eta_{\hat{\varepsilon}_n(x)}(x - y) |u(x) - u(y)| dy \right) g(x) dx \\ &\leq \int_0^1 \int_D \left(\frac{1}{\hat{\varepsilon}_n(x)} \int_{B(x, \hat{\varepsilon})} \eta_{\hat{\varepsilon}_n(x)}(x - y) |\nabla u(x + t(y - x)) \cdot (x - y)| dy \right) g(x) dx dt, \end{aligned}$$

which follows from the fundamental theorem of calculus. Now, for every fixed $x \in D$, we use the change of variables $h := \frac{y-x}{\hat{\varepsilon}_n(x)}$ in the above expression to rewrite

$$\frac{1}{\hat{\varepsilon}_n(x)} \int_{B(x, \hat{\varepsilon}_n(x))} \eta_{\hat{\varepsilon}_n(x)}(x - y) |\nabla u(x + t(y - x)) \cdot (x - y)| dy = \int_{B(0,1)} \eta(h) |\nabla u(x + t\hat{\varepsilon}_n(x)h) \cdot h| dh.$$

But because $u \in C_c^\infty(\mathbb{R}^d)$ and $\sup_{x \in D} \hat{\varepsilon}_n(x) \rightarrow 0$ as $n \rightarrow \infty$, it follows that

$$\lim_{n \rightarrow \infty} \sup_{x \in D} \left| \int_{B(0,1)} \eta(h) |\nabla u(x + t\hat{\varepsilon}_n(x)h) \cdot h| dh - \int_{B(0,1)} \eta(h) |\nabla u(x) \cdot h| dh \right| = 0.$$

Hence,

$$\limsup_{n \rightarrow \infty} E_n(u) \leq \int_0^1 \int_D \left(\int_{B(0,1)} \eta(h) |\nabla u(x) \cdot h| dh \right) g(x) dx dt = \sigma_\eta TV(u; g).$$

□

With the previous lemma at hand we can now establish Proposition 3.1.

Proof of Proposition 3.1. We make use of Lemma 3.2 and for that purpose we first need to approximate the function $\rho : D \rightarrow \mathbb{R}$ from above and from below using appropriate Lipschitz functions. More precisely, for every $s \in \mathbb{N}$ let

$$\tilde{\rho}_{1,s}(x) := \inf_{y \in D} \{\rho(y) + s|x - y|\}, \quad \rho_{2,s}(x) := \sup_{y \in D} \{\rho(y) - s|x - y|\}.$$

Notice that the functions $\tilde{\rho}_{1,s}, \rho_{2,s}$ are Lipschitz continuous, and as $s \rightarrow \infty$,

$$\rho_{1,s}(x) \nearrow \rho(x), \quad \rho_{2,s}(x) \searrow \rho(x), \quad \forall x \in D.$$

For every $s \in \mathbb{N}$ we modify the function $\tilde{\rho}_{1,s}$ slightly so as to create a Lipschitz function $\rho_{1,s}$ that in addition to minorizing ρ , is such that for all large enough $n \in \mathbb{N}$

$$(3.4) \quad \int_{B(x, \varepsilon_n(x))} \rho_{1,s}(y) dy \leq \int_{B(x, \varepsilon_n(x)) \cap D} \rho(y) dy = \nu(B(x, \varepsilon_n(x))), \quad \forall x \in D.$$

To achieve this, we use the regularity assumption on the boundary of D as follows. From the fact that D is an open and bounded set with Lipschitz boundary it follows (see [14], Theorem 1.2.2.2) that there exists a cone $\mathcal{C} \subseteq \mathbb{R}^d$ with non-empty interior and vertex at the origin, a family of rotations $\{R_x\}_{x \in D}$ and a number $1 > \zeta > 0$ such that for every $x \in D$,

$$x + R_x(\mathcal{C} \cap B(0, \zeta)) \subseteq D.$$

For every $s \in \mathbb{N}$, let $\xi_s : \mathbb{R}^d \rightarrow [0, 1]$ be a smooth cutoff function satisfying $\xi_s(x) \equiv 1$ in $\{y \in D : \text{dist}(y, \partial D) > 1/s\}$ and $\xi \equiv 0$ in $\{y \in D : \text{dist}(y, \partial D) < 1/2s\}$, and let us define

$$\rho_{1,s}(x) := \frac{|B(0, 1) \cap \mathcal{C}|}{|B(0, 1)|} \rho_{\min}(1 - \xi_s(x)) + \tilde{\rho}_{1,s}(x) \xi_s(x), \quad x \in \mathbb{R}^d.$$

It is clear that the functions $\rho_{1,s}$ are Lipschitz, they minorize ρ and they converge to ρ pointwise. In addition, for any fixed s , if $n \in \mathbb{N}$ is large enough so that in particular $\sup_{x \in D} \varepsilon_n(x) < \min\{\zeta, \frac{1}{8s}\}$, then (3.4) holds. Indeed, notice that if $x \in D$ is $1/4s$ units away from ∂D then (3.4) follows directly from the fact that $\rho_{1,s} \leq \rho$; on the other hand, if $x \in D$ is within distance $1/4s$ from ∂D , we conclude that

$$\begin{aligned} \int_{B(x, \hat{\varepsilon}_n(x))} \rho_{1,s}(y) dy &= |\mathcal{C} \cap B(0, 1)| \rho_{\min}(\hat{\varepsilon}_n(x))^d \leq \int_{(x + R_x(\mathcal{C})) \cap B(x, \hat{\varepsilon}_n(x))} \rho(y) dy \\ &\leq \int_{B(x, \hat{\varepsilon}_n(x)) \cap D} \rho(y) dy = \nu(B(x, \hat{\varepsilon}_n(x))). \end{aligned}$$

Let us now establish the limsup inequality. We introduce the functionals $F_{n,s} : L^1(D) \rightarrow [0, \infty)$ given by

$$F_{n,s}(u) := \int_D \tilde{f}_{n,s}(x) \left(\int_D \eta_{\varepsilon_n(x)}(x - y) |u(x) - u(y)| \rho_{2,s}(y) dy \right) \rho_{2,s}(x) dx, \quad u \in L^1(D),$$

where

$$\tilde{f}_{n,s}(x) := \frac{(\hat{\varepsilon}_n(x))^d}{(\mu_s(B(x, \varepsilon_n(x))))^{1+1/d}},$$

and

$$d\mu_s(x) := \rho_{1,s}(x)dx.$$

It follows that for every fixed $s \in \mathbb{N}$, for all large $n \in \mathbb{N}$

$$F_n(v) \leq F_{n,s}(v), \quad \forall v \in L^1(D).$$

Hence, from (3.3) it follows that for every arbitrary $u \in C_c^\infty(\mathbb{R}^d)$

$$\limsup_{n \rightarrow \infty} F_n(u) \leq \limsup_{n \rightarrow \infty} F_{n,s}(u) \leq \frac{\sigma_\eta}{\alpha_d^{1+1/d}} \int_D \frac{\rho_{2,s}^2(x)}{\rho_{1,s}^{1+1/d}(x)} d|Du|(x).$$

Taking $s \rightarrow \infty$ in the above inequality we deduce that

$$\limsup_{n \rightarrow \infty} F_n(u) \leq \frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(u; \rho^{1-1/d}).$$

Remark (2.14) and the density of $C_c^\infty(\mathbb{R}^d)$ functions with respect to TV imply the desired result.

To establish the liminf inequality, it is enough to change the roles of $\rho_{1,s}$ and $\rho_{2,s}$ in the definition of the functional $F_{n,s}$ and use Lemma 3.2 to conclude that if $u_n \xrightarrow{L^1(D)} u$, then for every $s \in \mathbb{N}$

$$\liminf_{n \rightarrow \infty} F_n(u_n) \geq \liminf_{n \rightarrow \infty} F_{n,s}(u_n) \geq \frac{\sigma_\eta}{\alpha_d^{1+1/d}} \int_D \frac{\rho_{1,s}^2(x)}{\rho_{2,s}^{1+1/d}(x)} d|Du|(x).$$

Taking the limit as $s \rightarrow \infty$ on the right hand side of the above expression we obtain the desired result. □

We are ready to establish Theorem 2.7.

Proof of liminf inequality in Theorem 2.7: Let us consider the cone $\mathcal{C}$, the family of rotations $\{R_x\}_{x \in D}$, and the number $1 > \zeta > 0$ introduced in the proof of Proposition 3.1. For $0 \leq \gamma_1 < \gamma_2 < \zeta$ and $x \in D$ let $A(x; \gamma_1, \gamma_2)$ be the annulus $A(x; \gamma_1, \gamma_2) := B(x, \gamma_2) \setminus B(x, \gamma_1)$. Then, there exist constants $\mathcal{K}_1$ (only depending on d , and on the cone $\mathcal{C}$) and $\mathcal{K}_2$ (only depending on d), such that for every $0 \leq \gamma_1 < \gamma_2 < \zeta$,

$$(3.5) \quad \mathcal{K}_1 \rho_{\min}(\gamma_2)^{d-1} \cdot (\gamma_2 - \gamma_1) \leq \nu(A(x; \gamma_1, \gamma_2) \cap (x + R_x(\mathcal{C}))) \leq \nu(A(x; \gamma_1, \gamma_2)), \quad \forall x \in D,$$

and

$$(3.6) \quad \nu(A(x; \gamma_1, \gamma_2)) \leq \mathcal{K}_2 \rho_{\max}(\gamma_2)^{d-1} \cdot (\gamma_2 - \gamma_1), \quad \forall x \in D.$$

We work in a set with full probability for which the maps $\{T_n\}_{n \in \mathbb{N}}$ from (2.7) exist. For every $x \in D$ we let $\varepsilon_n(x)$ be the number for which

$$\nu(B(x, \varepsilon_n(x))) = \bar{\varepsilon}_n^d = \frac{k_n}{n}.$$

On the other hand, we define $\dot{\varepsilon}_n(x)$ and $\hat{\varepsilon}_n(x)$ to be

$$\dot{\varepsilon}_n(x) := \varepsilon_n(x) - \frac{\mathcal{K}_2 \rho_{\max} \delta_n}{\mathcal{K}_1 \rho_{\min}},$$

$$\hat{\varepsilon}_n(x) := \dot{\varepsilon}_n(x) - 3\delta_n,$$

where δ_n is as in (2.7).

Notice that the assumption $n \gg k_n \gg \log(n)$ is equivalent to $1 \gg \bar{\varepsilon}_n \gg \delta_n$ where we recall $\bar{\varepsilon}_n$ is defined in (2.2). Combining this with (3.5) we deduce that

$$(3.7) \quad \lim_{n \rightarrow \infty} \sup_{x \in D} \varepsilon_n(x) = 0, \quad \lim_{n \rightarrow \infty} \sup_{x \in D} \left| \frac{\varepsilon_n(x)}{\hat{\varepsilon}_n(x)} - 1 \right| = 0, \quad \lim_{n \rightarrow \infty} \frac{\delta_n}{\inf_{x \in D} \hat{\varepsilon}_n(x)} = 0.$$

Now, from (3.5) we see that for all large enough n , and for all $x \in D$,

$$(3.8) \quad \begin{aligned} \nu(B(x, \dot{\varepsilon}_n(x))) &= \nu(B(x, \varepsilon_n(x))) - \nu(A(x; \dot{\varepsilon}_n(x), \varepsilon_n(x))) \\ &\leq \frac{k_n}{n} - \mathcal{K}_2 \rho_{\max}(\varepsilon_n(x))^{d-1} \delta_n. \end{aligned}$$

Thus,

$$\begin{aligned} \nu_n(B(x, \dot{\varepsilon}_n(x))) &= \nu_n(B(x, \dot{\varepsilon}_n(x))) - \nu(B(x, \dot{\varepsilon}_n(x))) + \nu(B(x, \dot{\varepsilon}_n(x))) \\ &\leq \nu(B(x, \dot{\varepsilon}_n(x) + \|Id - T_n\|_\infty)) - \nu(B(x, \dot{\varepsilon}_n(x))) + \nu(B(x, \dot{\varepsilon}_n(x))) \\ &\leq \mathcal{K}_2 \rho_{\max} \varepsilon_n(x)^{d-1} \|Id - T_n\|_\infty + \frac{k_n}{n} - \mathcal{K}_2 \rho_{\max}(\varepsilon_n(x))^{d-1} \cdot \delta_n \\ &\leq \mathcal{K}_2 \rho_{\max} \varepsilon_n(x)^{d-1} \delta_n + \frac{k_n}{n} - \mathcal{K}_2 \rho_{\max} \varepsilon_n(x)^{d-1} \delta_n \\ &= \frac{k_n}{n}, \end{aligned}$$

where in the first inequality we have used the fact that T_n is a transportation map between ν and ν_n , and in the second inequality we have used (3.6). Combining the previous inequality with the fact that $B(T_n(x), \dot{\varepsilon}_n(x) - \delta_n) \subseteq B(x, \dot{\varepsilon}_n(x))$ we deduce that

$$\nu_n(B(T_n(x), \dot{\varepsilon}_n(x) - \delta_n)) \leq \frac{k_n}{n}.$$

In particular, if $y \in D$ is such that $T_n(y) \not\prec_{k_n} T_n(x)$, then from Remark 2.1 it follows that

$$\hat{\varepsilon}_n(x) = \dot{\varepsilon}_n(x) - 3\delta_n < |T_n(x) - T_n(y)| - 2\delta_n \leq |T_n(x) - T_n(y)| - |x - T_n(x)| - |y - T_n(y)| \leq |x - y|.$$

We conclude that for all large enough n , $J_{k_n}(T_n(x), T_n(y)) = 0$ implies $\hat{\varepsilon}_n(x) < |x - y|$. In particular, we deduce that for all large enough $n \in \mathbb{N}$,

$$(3.9) \quad J_{k_n}(T_n(x), T_n(y)) \geq \eta \left(\frac{|x - y|}{\hat{\varepsilon}_n(x)} \right), \quad \forall x, y \in D.$$

Summarizing, by contracting the radius $\varepsilon_n(x)$ a little bit we obtained $\hat{\varepsilon}_n(x)$ satisfying (3.7) and (3.9).

Let us now consider a sequence of functions $\{u_n\}_{n \in \mathbb{N}}$ with $u_n \in L^1(\nu_n)$ for which $u_n \xrightarrow{TL^1} u$ for some $u \in L^1(\nu)$. It follows from (3.9) that for all large enough $n \in \mathbb{N}$,

$$(3.10) \quad \begin{aligned} GTV_{n, k_n}(u_n) &= \frac{1}{n^2 \bar{\varepsilon}_n^{d+1}} \sum_{i, j} J_{k_n}(\mathbf{x}_i, \mathbf{x}_j) |u_n(\mathbf{x}_i) - u_n(\mathbf{x}_j)| \\ &= \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\bar{\varepsilon}_n^{d+1}} \left(\sum_{j=1}^n J_{k_n}(\mathbf{x}_i, \mathbf{x}_j) |u_n(\mathbf{x}_i) - u_n(\mathbf{x}_j)| \right) \\ &= \int_D \frac{1}{\bar{\varepsilon}_n^{d+1}} \left(\int_D J_{k_n}(T_n(x), T_n(y)) |u_n \circ T_n(x) - u_n \circ T_n(y)| d\nu(y) \right) d\nu(x) \\ &\geq \int_D \frac{1}{\bar{\varepsilon}_n^{d+1}} \left(\int_D \eta \left(\frac{|x - y|}{\hat{\varepsilon}_n(x)} \right) |u_n \circ T_n(x) - u_n \circ T_n(y)| d\nu(y) \right) d\nu(x) \\ &= \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(|x - y|) |u_n \circ T_n(x) - u_n \circ T_n(y)| d\nu(y) \right) d\nu(x) \\ &= F_n(u_n \circ T_n), \end{aligned}$$

where for an arbitrary function $v \in L^1(\nu)$, $F_n(v)$ is defined as

$$F_n(v) := \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(|x - y|) |v(x) - v(y)| d\nu(y) \right) d\nu(x),$$

and the function $\hat{f}_n$ is given by $\hat{f}_n(x) := \frac{\hat{\varepsilon}_n(x)^d}{(\nu(B(x, \varepsilon_n(x))))^{1+1/d}}$. Since $u_n \xrightarrow{TL^1} u$ implies that $u_n \circ T_n \xrightarrow{L^1(\nu)} u$, it follows from Proposition (3.1) and (3.10) that

$$\liminf_{n \rightarrow \infty} GTV_{n,k_n}(u_n) \geq \liminf_{n \rightarrow \infty} F_n(u_n \circ T_n) \geq \frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(u; \rho^{1-1/d}),$$

which establishes the liminf inequality in Theorem 2.7. $\square$

Proof of the limsup inequality in Theorem (2.7): We work in a set with full probability for which the maps $\{T_n\}_{n \in \mathbb{N}}$ from (2.7) exist. By an analogue construction to the one used in the proof of the liminf inequality (this time enlarging $\varepsilon_n(x)$ instead of contracting it), we may construct a function $\hat{\varepsilon}_n$ satisfying (3.7) and

$$J_{k_n}(T_n(x), T_n(y)) \leq \eta \left(\frac{|x - y|}{\hat{\varepsilon}_n(x)} \right), \quad \forall x, y \in D.$$

Let $u \in C_c^\infty(\mathbb{R}^d)$ and for every $n \in \mathbb{N}$ let u_n be the function in $L^1(\nu_n)$ defined by

$$u_n(\mathbf{x}_i) := u(\mathbf{x}_i), \quad i = 1, \dots, n.$$

In other words, u_n is the function u restricted to the point cloud X_n . Then,

$$\begin{aligned} GTV_{n,k_n}(u_n) &= \frac{1}{n^2 \bar{\varepsilon}_n^{d+1}} \sum_{i,j} J(\mathbf{x}_i, \mathbf{x}_j) |u(\mathbf{x}_i) - u(\mathbf{x}_j)| \\ &= \int_D \frac{1}{\bar{\varepsilon}_n^{d+1}} \left(\int_D J(T_n(x), T_n(y)) |u \circ T_n(x) - u \circ T_n(y)| d\nu(y) \right) d\nu(x) \\ (3.11) \quad &\leq \int_D \frac{1}{\bar{\varepsilon}_n^{d+1}} \left(\int_D \eta \left(\frac{|x - y|}{\hat{\varepsilon}_n(x)} \right) |u_n \circ T_n(x) - u_n \circ T_n(y)| d\nu(y) \right) d\nu(x) \\ &\leq \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(|x - y|) |u(x) - u(y)| d\nu(y) \right) d\nu(x) + \mathcal{I}_n + \mathcal{II}_n \\ &= F_n(u) + \mathcal{I}_n + \mathcal{II}_n, \end{aligned}$$

where

$$\mathcal{I}_n := \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(|x - y|) |u(x) - u(T_n(x))| d\nu(y) \right) d\nu(x),$$

and

$$\mathcal{II}_n := \int_D \hat{f}_n(x) \left(\int_D \eta_{\hat{\varepsilon}_n(x)}(|x - y|) |u(y) - u(T_n(y))| d\nu(y) \right) d\nu(x),$$

But notice that

$$\mathcal{I}_n, \mathcal{II}_n \leq \frac{C \|\nabla u\|_\infty \|Id - T_n\|_\infty}{\inf_{x \in D} \hat{\varepsilon}_n(x)},$$

for some constant C . Hence, $\lim_{n \rightarrow \infty} \mathcal{I}_n = \lim_{n \rightarrow \infty} \mathcal{II}_n = 0$. It follows from the proof of Proposition (3.1) that

$$\limsup_{n \rightarrow \infty} GTV_{n,k_n}(u_n) \leq \limsup_{n \rightarrow \infty} F_n(u) \leq \frac{\sigma_\eta}{\alpha_d^{1+1/d}} TV(u; \rho^{1-1/d}).$$

Using the density of $C_c^\infty(\mathbb{R}^d)$ with respect to TV , we may find a recovery sequence $\{u_n\}_{n \in \mathbb{N}}$ with $u_n \in L^1(\nu_n)$ for every $u \in L^1(D)$. Finally, if $u \in L^1(D)$ is of the form $u = \mathbf{1}_A$ for some measurable set $A \subseteq D$, we can actually choose this recovery sequence to consist of indicator functions, as it follows from the fact that the energies GTV_{n,k_n} satisfy a coarea formula (see the proof of Corollary 1.3 in [9]). This establishes the last statement in Theorem 2.7. $\square$

3.1. Proof of compactness.

Proof of Theorem 2.8. The compactness in Theorem 2.8 follows directly from the compactness result in [9]. Indeed, consider $\hat{\varepsilon}_n$ as defined in the proof of the liminf inequality of Theorem 2.7 and define

$$\varepsilon_n^l := \inf_{x \in D} \hat{\varepsilon}_n(x) - 2\delta_n.$$

Then it follows that for all large enough n ,

$$\frac{\varepsilon_n^l}{\bar{\varepsilon}_n} \geq c > 0,$$

for some constant $c > 0$. Moreover, for every i, j we have

$$(3.12) \quad J_{k_n}(\mathbf{x}_i, \mathbf{x}_j) \geq \eta \left(\frac{|\mathbf{x}_i - \mathbf{x}_j|}{\varepsilon_n^l} \right).$$

Let $\{u_n\}_{n \in \mathbb{N}}$ be a sequence with $u_n \in L^1(\nu_n)$ satisfying

$$\sup_{n \in \mathbb{N}} \|u_n\|_{L^1(\nu_n)} < \infty,$$

and

$$\sup_{n \in \mathbb{N}} GTV_{n, k_n}(u_n) < \infty.$$

It follows from inequality (3.12) that

$$GTV_{n, k_n}(u_n) \geq \frac{c^d}{n^2 \varepsilon_n^l} \sum_{i, j} \eta_{\varepsilon_n^l}(|\mathbf{x}_i - \mathbf{x}_j|) |u_n(\mathbf{x}_i) - u_n(\mathbf{x}_j)| =: c^d GTV_{n, \varepsilon_n^l}(u_n).$$

Therefore,

$$\sup_{n \in \mathbb{N}} GTV_{n, \varepsilon_n^l}(u_n) < \infty.$$

The pre-compactness of $\{u_n\}_{n \in \mathbb{N}}$ in TL^1 follows from Theorem 1.2 in [9]. $\square$

REFERENCES

- [1] E. ARIAS-CASTRO, B. PELLETIER, AND P. PUDLO, *The normalized graph cut and Cheeger constant: from discrete to continuous*, Adv. in Appl. Probab., 44 (2012), pp. 907–937.
- [2] A. BALDI, *Weighted BV functions*, Houston J. Math., 27 (2001), pp. 683–705.
- [3] M. BELKIN AND P. NIYOGI, *Convergence of Laplacian eigenmaps*, Advances in Neural Information Processing Systems (NIPS), 19 (2007), p. 129.
- [4] ———, *Towards a theoretical foundation for Laplacian-based manifold methods*, J. Comput. System Sci., 74 (2008), pp. 1289–1308.
- [5] X. BRESSON, T. LAURENT, D. UMINSKY, AND J. VON BRECHT, *Multiclass total variation clustering*, in Advances in Neural Information Processing Systems 26, C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K. Weinberger, eds., 2013, pp. 1421–1429.
- [6] R. R. COIFMAN AND S. LAFON, *Diffusion maps*, Appl. Comput. Harmon. Anal., 21 (2006), pp. 5–30.
- [7] G. DAL MASO, *An Introduction to Γ -convergence*, Springer, 1993.
- [8] N. GARCÍA TRILLOS AND R. MURRAY, *A new analytical approach to consistency and overfitting in regularized empirical risk minimization*. Preprint, 2016.
- [9] N. GARCÍA TRILLOS AND D. SLEPČEV, *Continuum limit of total variation on point clouds*, Archive for Rational Mechanics and Analysis, (2015), pp. 1–49.
- [10] N. GARCÍA TRILLOS AND D. SLEPČEV, *On the rate of convergence of empirical measures in ∞ -transportation distance*, Canad. J. Math., 67 (2015), pp. 1358–1383.
- [11] N. GARCÍA TRILLOS AND D. SLEPČEV, *A variational approach to the consistency of spectral clustering*. Preprint., 2015.
- [12] N. GARCÍA TRILLOS, D. SLEPČEV, J. VON BRECHT, T. LAURENT, AND X. BRESSON, *Consistency of Cheeger and ratio graph cuts*. to appear in Journal of Machine Learning Research, 2015.
- [13] E. GINÉ AND V. KOLTCHINSKII, *Empirical graph Laplacian approximation of Laplace-Beltrami operators: large sample results*, in High dimensional probability, vol. 51 of IMS Lecture Notes Monogr. Ser., Inst. Math. Statist., Beachwood, OH, 2006, pp. 238–259.
- [14] P. GRISVARD, *Elliptic problems in nonsmooth domains*, vol. 24 of Monographs and Studies in Mathematics, Pitman (Advanced Publishing Program), Boston, MA, 1985.

- [15] M. HEIN, J.-Y. AUDIBERT, AND U. VON LUXBURG, *From graphs to manifolds—weak and strong pointwise consistency of graph Laplacians*, in Learning theory, Springer, 2005, pp. 470–485.
- [16] M. HEIN AND T. BÜHLER, *An inverse power method for nonlinear eigenproblems with applications in 1-spectral clustering and sparse PCA*, in Advances in Neural Information Processing Systems (NIPS), 2010, pp. 847–855.
- [17] M. HEIN AND S. SETZER, *Beyond spectral clustering - tight relaxations of balanced graph cuts*, in Advances in Neural Information Processing Systems (NIPS), J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K. Weinberger, eds., 2011, pp. 2366–2374.
- [18] G. LEONI, *A first course in Sobolev spaces*, vol. 105 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2009.
- [19] F. MAGGI, *Sets of finite perimeter and geometric variational problems*, vol. 135 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.
- [20] A. C. PONCE, *A new approach to Sobolev spaces and connections to Γ -convergence*, Calc. Var. Partial Differential Equations, 19 (2004), pp. 229–255.
- [21] S. S. RANGAPURAM AND M. HEIN, *Constrained 1-spectral clustering*, in International conference on Artificial Intelligence and Statistics (AISTATS), 2012, pp. 1143–1151.
- [22] J. SHI AND J. MALIK, *Normalized cuts and image segmentation*, Pattern Analysis and Machine Intelligence, IEEE Transactions on, 22 (2000), pp. 888–905.
- [23] A. SINGER, *From graph to manifold Laplacian: the convergence rate*, Appl. Comput. Harmon. Anal., 21 (2006), pp. 128–134.
- [24] A. SINGER AND H.-T. WU, *Spectral convergence of the connection laplacian from random samples*, 2015.
- [25] D. TING, L. HUANG, AND M. I. JORDAN, *An analysis of the convergence of graph Laplacians*, in Proceedings of the 27th International Conference on Machine Learning, 2010.
- [26] U. VON LUXBURG, *A tutorial on spectral clustering*, Statistics and Computing, 17 (2007), pp. 395–416.
- [27] U. VON LUXBURG, M. BELKIN, AND O. BOUSQUET, *Consistency of spectral clustering*, Ann. Statist., 36 (2008), pp. 555–586.

DIVISION OF APPLIED MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RI, 02912, USA.

Email: nicolas_garcia_trillos@brown.edu